

Tree stem bases are sources of CH₄ and N₂O in a tropical forest on upland soil during the dry to wet season transition

Welch, Bertie; Gauci, Vincent; Sayer, Emma J.

DOI:

[10.1111/gcb.14498](https://doi.org/10.1111/gcb.14498)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Welch, B, Gauci, V & Sayer, EJ 2019, 'Tree stem bases are sources of CH₄ and N₂O in a tropical forest on upland soil during the dry to wet season transition', *Global Change Biology*, vol. 25, no. 1, pp. 361-372.
<https://doi.org/10.1111/gcb.14498>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Tree stem bases are sources of CH₄ and N₂O in a tropical forest on upland soils during the dry to wet season transition

Authors: Bertie Welch¹, Vincent Gauci¹ & Emma J. Sayer^{2,3}

Addresses:

1. School of Environment, Earth & Ecosystem Science, The Open University, Milton Keynes, MK7 6AA, United Kingdom

2. Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom

3. Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancón, Republic of Panama.

Corresponding author:

E.J. Sayer

e.sayer@lancaster.ac.uk

Keywords: Methane, nitrous oxide, tree stem emissions, upland tropical forest, leaf litter, trace greenhouse gases, soil

Type: Primary Research Article

Abstract

Tropical forests on upland soils are assumed to be a methane (CH₄) sink and a weak source of nitrous oxide (N₂O), but studies of wetland forests have demonstrated that tree stems can be a substantial source of CH₄, and recent evidence from temperate woodlands suggests that tree stems can also emit N₂O. Here, we measured CH₄ and N₂O fluxes from the soil and from tree stems in a semi-evergreen tropical forest on upland soil. To examine the influence of seasonality, soil abiotic conditions, and substrate availability (litter inputs) on trace greenhouse gas (GHG) fluxes, we conducted our study during the transition from the dry to the wet season in a long-term litter manipulation experiment in Panama, Central America. Trace GHG fluxes were measured from individual stem bases of two common tree species and from soils beneath the same trees. Soil CH₄ fluxes varied from uptake in the dry season to minor emissions in the wet season. Soil N₂O fluxes were negligible during the dry season but increased markedly after the start of the wet season. By contrast, tree stem bases emitted CH₄ and N₂O throughout the study. Although we observed no clear effect of litter manipulation on trace GHG fluxes, tree species and litter treatments interacted to influence CH₄ fluxes from stems and N₂O fluxes from stems and soil, indicating complex relationships between tree species traits and decomposition processes that can influence trace GHG dynamics. Collectively, our results show that tropical trees can act as conduits for trace GHGs that most likely originate from deeper soil horizons, even when they are growing on upland soils. Coupled with the finding that the soils may be a weaker sink for CH₄ than previously thought, our research highlights the need to reappraise trace gas budgets in tropical forests.

Introduction

Methane (CH_4) and nitrous oxide (N_2O) are important trace greenhouse gases (GHGs) with radiative effects 25 and 298 times greater than CO_2 , respectively (Houghton et al., 2001). Interest in trace GHG exchange in tropical forests has grown in recent years, particularly in saturated wetland areas of the tropics such as the Amazon floodplain (Graffman et al., 2008; Pangala et al., 2017) and mangrove swamps (Kreuzwieser et al., 2003; Krithika et al., 2008). Natural wetlands are the single largest individual source of atmospheric methane contributing 177-284 Tg CH_4 yr^{-1} (IPCC, 2013), to which tropical wetland emissions from a variety of sources (including waterlogged soils) are a significant contributor, but the contribution of tropical forests on upland soils (i.e. soils that are rarely flooded and only temporarily water-saturated) has not yet been quantified. Globally, emissions of nitrous oxide from soils in natural ecosystems account for 37% of total global surface emissions (IPCC, 2007), estimated at 3.37-6.60 Tg N yr^{-1} (Zhuang et al., 2012) and tropical rainforests are the single biggest natural source of N_2O (Bouwman et al., 1995).

Tree stems can also emit significant amounts of CH_4 in temperate (Gauci et al., 2010) and tropical (Pangala et al., 2013; Pangala et al., 2017) wetland ecosystems, and mesocosm experiments showed that black alder trees, typical of European temperate wetlands, can also act as a pathway for N_2O emissions to the atmosphere (Rusch and Rennenberg, 1998). CH_4 and N_2O are produced under anoxic conditions in waterlogged soils by methanogenic consortia of archaea or denitrifying bacteria, respectively, and the gases diffuse into soil water, and then from water into roots as either a solute or a gas. The gases move up the tree stem by either mass flow (transpiration or pressurized ventilation) or diffusion, then diffuse from the xylem through the stem tissue to the atmosphere via lenticels and other structures that aid gas exchange (Carmichael et al., 2014). Findings extrapolated from glasshouse experiments suggest that wetland hardwood trees could account for emissions of around 60 Tg CH_4 yr^{-1} (Rice et al.,

2010); tree stem fluxes accounted for 62-87% of total ecosystem CH₄ flux in tropical peat forests in Indonesia (Pangala et al. 2013) and contribute half of all emitted methane in the Amazon floodplain (~20 Tg; Pangala et al., 2017).

We are only just beginning to understand the role of tropical tree stems as conduits of soil-produced trace GHGs, and the vast majority of research on this subject to date has been in forested wetlands. Wetland tree species have evolved a variety of specialist tissues to aid oxygen transport to roots in anoxic soils such as aerenchyma, increased stem lenticel density (Pangala et al., 2014) and adventitious roots, which can transport soil-generated CH₄ in mangrove tree species (Purvaja et al., 2004). Inter-species variations in wood specific density and stem lenticel numbers could be important controls of stem emissions, as tree stem CH₄ flux is negatively related to wood density and positively related to lenticel number (Pangala et al., 2013). Tree stem emissions of trace GHGs can occur on upland soils when such soils become saturated with water, which reduces soil oxygen concentrations and facilitates the activity of anoxic methanogenic archaea and denitrifying bacteria that thrive in such soil conditions. Tree stem emissions of CH₄ and N₂O have been observed in temperate trees that lack aerenchyma and other adaptations to wet, anoxic soil conditions (Machacova *et al.*, 2013, 2016; Wen et al., 2017). In addition, recent work demonstrates production of CH₄ within tree stems (Wang et al., 2016) and high abundance of methanogens in heartwood (Yip et al., 2018). However, despite the global importance of tropical forests in GHG budgets, we know very little about CH₄ emissions from tropical tree stems on upland soils and we do not know whether they are also capable of emitting N₂O.

Although tropical forests on upland soils cover a larger land-surface area than tropical wetlands (Pan et al., 2013), they are generally not considered to be a major source of CH₄ and N₂O emissions. However, the role of tree stems as conduits of trace GHGs in tropical forests on upland soils warrants further attention because many tropical tree species have large buttress

94 roots and the greatest stem gas emissions are measured within 0.3-m of the soil surface (Rusch
95 and Rennenberg, 1998; Gauci et al., 2010; Pangala et al., 2013). Hence, even minor trace GHG
96 emissions from tropical trees on upland soils could represent a significant source of CH₄ and
97 N₂O. Indeed, recent work in temperate woodland demonstrated that tree stem emissions
98 diminish the methane sink of upland forests (Pitz & Megonigal, 2017), if the same applies to
99 tropical forests, it would affect global trace GHG budgets.

100 Regardless of whether soils are waterlogged or well-drained, the production of trace GHGs
101 in soils depends on the availability of suitable substrates (Li et al., 2000). Litter quantity can
102 influence the rates of trace GHG emissions from forest soils as it provides substrate (acetate
103 and hydrogen) for methanogens and the nitrate used in denitrification (Teh et al., 2008). The
104 potential link between litter inputs and trace GHG emissions from soil was explored by a study
105 of soil N₂O emissions from a wet forest in Costa Rica, in which doubling leaf litter inputs
106 increased rates of N₂O emissions by 43% relative to controls, with a corresponding decline of
107 42% in litter removal plots (Wieder et al., 2011). However, litter manipulation treatments in
108 subtropical forest in Southern China had no significant effect on soil CH₄ uptake or N₂O
109 production, implying that abiotic conditions in the mineral soil may be more important than
110 litter quantity (Tang et al., 2006). The effects of litter manipulation on trace GHG emissions
111 from tree stems is presently unknown, but as changes in mineral soil chemistry from litter were
112 the primary driver of changes in CH₄ and N₂O fluxes (Fender et al., 2013), it is conceivable that
113 litter manipulation could also affect tree stem emissions.

114 Hence, although trees can be a major conduit for CH₄ in tropical floodplains and peat forests,
115 we know very little about tree stem fluxes of CH₄ in tropical forests on upland soils, and there
116 are no field data on tree stem N₂O emissions in the tropics. We aimed to address these gaps in
117 our knowledge of tropical GHG emissions by measuring CH₄ and N₂O fluxes from the soil and
118 tree stems in a seasonal moist tropical forest on upland soils. We focussed our attention on

quantifying seasonal changes in CH₄ and N₂O fluxes at the base of tree stems, and we assessed the specific role of litter in providing substrate for trace GHG production. Accordingly, we conducted our study in an existing long-term litter manipulation experiment during the transition from the dry season to the wet season to test the following hypotheses:

- 1) Tree stems in tropical forests on upland soils will act as a conduit for CH₄ and N₂O produced in the soil; patterns in trace GHG emissions from tree stems will therefore mimic those from the soil, increasing in the wet season and decreasing in the dry season.
- 2) Litter manipulation treatments alter substrate availability to microorganisms and will therefore influence CH₄ and N₂O fluxes from soils and tree stems; hence, trace GHG emissions will be greater in litter addition treatments and lower in litter removal plots relative to controls.

Materials and Methods

Field area and sampling

The study was carried out within the Gigante Litter Manipulation Project, approximately 5 km south of Barro Colorado Island (BCI) in Panama, Central America (Sayer & Tanner, 2010). The 15 plots were set up between 2000 and 2002; each plot measures 45-m × 45-m and the edges of the plots were trenched to a depth of 0.5-m, lined with plastic and then backfilled. Starting in January 2003, the litter is raked up and removed from five plots every month (L-) and added to five plots where it is spread as evenly as possible (L+); five plots were left as controls (CT; see Sayer et al. 2006 and Sayer & Tanner 2010 for a full description). The mean annual temperature at the weather station on nearby Barro Colorado Island (within 2 km of the study site) is 26°C, mean annual rainfall is 2,600 mm and there is a strong dry season from mid-December to mid-April (Leigh, 1999). During the study period, maximum and minimum air temperatures were 32.4°C and 24.3°C respectively, soil temperature ranged from 24.9 – 29.2°C

and soil water content (SWC) was between 14% and 40%. The soil in the plots is characterised as a moderately acidic Oxisol (Cavelier, 1992).

Two tree species were selected for this study: the fast-growing canopy tree *Simarouba amara* (Aubl.) and the shade-tolerant subcanopy tree *Heisteria concinna* (Standl.), which occur frequently throughout the study forest and are among the most abundant tree species in the experimental plots (12% of all trees with dbh >10 cm; Sayer and Tanner, 2010). Both species have relatively smooth bark and straight stems, which facilitated sampling and the species have distinct specific wood densities (0.38 g cm^{-3} for *Simarouba* and 0.64 g cm^{-3} for *Heisteria*; Condit et al., 2013), which is likely to influence trace GHG fluxes from stems (Pangala et al., 2013). Trees were mapped and marked using handheld GPS. One individual per species was chosen per plot but only 13 of the 15 experimental plots contained live mature individuals of *Simarouba*; hence the present study included trees in four plots per treatment, making 12 *Heisteria* and 12 *Simarouba* trees in total.

Greenhouse gas fluxes from the soil were measured using permanently installed soil collars located 2-3 m to the north and south of each tree. The collars were made from 120-mm long sections of polyvinyl chloride pipe (internal diameter: 200 mm), which were embedded into the soil to 30-mm depth. All collars were installed at least two weeks prior to sampling in March 2014 and an appropriate amount of litter was placed into the collars in the CT and L+ plots to achieve consistency with the surrounding forest floor. To determine CH_4 and N_2O emissions from the soil, a PVC lid with an inner seal of gas-tight neoprene foam was placed on top of the collar; a 15-ml air sample was taken by syringe via a septum in the lid immediately after closure and then again after 3, 6 and 10 minutes. Each sample was injected into pre-evacuated 12-ml borosilicate vials (ExetainerTM, LabCo Ltd, High Wycombe, UK). The suction when removing the lid after sampling demonstrated the integrity of the seal on the soil chamber. Soil

temperature at 0-6 cm depth was recorded adjacent to the collars using a Themapen (ETI Ltd, Worthing, UK).

Tree stem gas fluxes were measured using a flexible chamber made from a 450-mm × 300-mm sheet of polycarbonate, lined with neoprene foam (19 mm wide, 25 mm thick; Siegenthaler et al. 2016). The chambers were secured to the tree stems at 0.3-m above the forest floor using cam buckle straps. Gas samples were taken by syringe from a septum in the middle of the chamber at 0, 5, 10 and 15 minutes, and injected into pre-evacuated 12-ml vials as described above.

Air samples from the tree stem and soil chambers were collected every two weeks between 30 March and 20 July 2014. Air pressure and temperature outside the stem chamber were recorded at the start of sampling using a Thermometer-Hygrometer-Barometer probe (Commeter C4141; Comet Systems, Czech Republic). Soil temperature at 0-10-cm depth was measured adjacent to the collars using a soil temperature probe and volumetric soil water content at 0-6 cm depth was measured using a Thetaprobe (Delta-T Devices, Cambridge, UK) calibrated to local soil conditions following the manufacturer's instructions. Collection of soil temperature and soil water content data during gas sampling was limited to 28 May - 14 June 2014 and 2 - 6 July 2014 due to equipment malfunction. We therefore used monthly values measured in the plots as part of a separate study (Sayer et al. unpublished data) and weekly rainfall data measured at the meteorological tower on Barro Colorado Island (courtesy of the Physical Monitoring Program of the Smithsonian Tropical Research Institute).

The CH₄ concentrations of the air samples were analysed within a week of sampling using off-axis Integrated Cavity Output Spectroscopy (FMA-200 Fast Methane Analyser; Los Gatos Research, Mountain View, CA, USA). The N₂O concentrations of the air samples were analysed using gas chromatography (Ai 94 Gas Chromatograph, Cambridge Instruments, Ellutia, Ely, UK). All methods for the measurement of GHGs, including the testing of the

chamber method and sensitivity of measurements are discussed fully in Siegenthaler et al. (2016).

Data analyses

The data were inspected visually before further analysis; we considered extreme outliers that lay outside of the 5th - 95th interquartile range as measurement errors and removed them from the dataset. For soil fluxes, we omitted two data-points for CH₄ and one value for N₂O (out of 188 and 108, respectively) and for tree stem fluxes, we omitted 20 data-points for CH₄ and three data-points for N₂O (out of 201 and 115, respectively).

All data analyses were conducted in R 3.3.2 (R Core Team, 2016) using the lme4 package for mixed effects models (Bates et al., 2015). Gas fluxes were calculated for each chamber following Baird et al. (2010), whereby the least squares linear regression slope of the four sample concentrations is plotted against sampling time and the slope to give the gas flux in $\mu\text{g m}^{-2} \text{ h}^{-1}$. Gas flux measurements were only used for further statistical analysis if the R^2 of the regression was >0.7 ; this cut-off point was chosen following Alm et al. (2007; cited in Cooper et al., 2014), who noted that low fluxes (especially those near to zero) tend to have low R^2 values. Concentration changes in N₂O in dry season samples were too small to estimate non-zero fluxes (i.e. $R^2 < 0.7$); we therefore only present wet season data for soil and tree stem N₂O fluxes.

Soil water content, soil temperature, and air temperature were strongly correlated; however, as is typical of the tropics, temperature only varied within a narrow range. Given that rainfall and soil moisture exhibited far larger variation (and have a fundamental control on soil CH₄ and N₂O production), we investigated the relationships between soil water content and trace GHG fluxes from soils and tree stems using monthly means. Relationships were inspected visually and emerging patterns were then tested using linear models. We then assessed the

218 influence of tree species, litter treatment, and their interaction on soil and stem trace GHG fluxes
 219 using linear mixed effects models (*lmer* function) with plot and time as random effects. The
 220 significance of each term was determined by comparing nested models using likelihood ratio
 221 tests. Models were simplified by sequentially dropping terms until a minimum adequate model
 222 was reached, using AICs and p-values to check for model improvement (Pinheiro and Bates,
 223 2000). The final model fit was inspected using diagnostic plots. Effects of seasonal variation
 224 were tested by comparing minimum adequate models with and without time as a random effect.
 225 Statistics for mixed effects models are given for the comparison between the best-fit model and
 226 the corresponding null model. All results are reported as significant at $p < 0.05$ but due to the
 227 low number of replicate plots ($n = 4$), marginally significant trends are also reported at $p < 0.1$.




Figure 1. The relationship between methane fluxes from the soil and soil water content at 0-6 cm depth in a lowland tropical forest on upland soil in Panama, Central America, during the transition from the dry season (circles) to the wet season (triangles) from March to July 2014.

228

229 Results

230 Seasonal variation in CH_4 fluxes

Soil water content was strongly related to total rainfall ($R^2 = 0.6$, $p < 0.01$) and CH₄ fluxes increased with soil water content ($R^2 = 0.3$, $p < 0.01$; Fig. 1). Soil CH₄ fluxes therefore varied significantly between the dry and wet season ($p < 0.001$, $\chi^2 = 36.4$), whereby soils acted as a CH₄ sink during the dry season and switched to being a source within two to three weeks of the start of the wet season (Fig. 2a).

There was no clear seasonal pattern for tree stem CH₄ fluxes; although stem CH₄ emissions tended to be larger during the wet season, they were not significantly so (Fig. 2a) and there were no significant relationships between tree stem fluxes and soil water content.




Figure 2 Methane (CH_4) fluxes from tree stems at 0.3-m height (top panel) and soil (centre panel) in a lowland tropical forest on upland soil in Panama, Central America; data shown are pooled across experimental plots with three litter manipulation treatments, means and standard errors are therefore given for $n = 12$; measurements were made weekly during the transition from the dry season to the wet season (bottom panel, weeks 14-19 and 20-30, respectively), with corresponding changes in total rainfall (bars) and temperature (line).

240 *Species and treatment effects on soil CH₄ fluxes*

241 There were no effects of species, treatment or their interaction on soil CH₄ fluxes. Soil CH₄
 242 fluxes remained predominantly negative until week 24 under individuals of *Heisteria* and until
 243 week 21 under individuals of *Simarouba*, indicating dry season uptake of CH₄ before a
 244 transition to emissions within two to four weeks after the first heavy rainfall of the year (Figs.
 245 2b and 3). The median flux beneath *Heisteria* was 8.3 $\mu\text{g m}^{-2} \text{hr}^{-1}$, which was slightly higher
 246 than the median CH₄ flux of 6.3 $\mu\text{g m}^{-2} \text{hr}^{-1}$ from chambers under *Simarouba*.

247




Figure 3. Soil methane (CH₄) fluxes under individuals of two common tree species: *Heisteria concinna* (top panels) and *Simarouba amara* (bottom panels) in experimental litter removal (L-), control (CT) and litter addition (L+) treatments in a lowland tropical forest on upland soil in Panama, Central America, measured weekly during the transition from the dry season (weeks 14-19) to the wet season (weeks 20-30); ranges (boxes and whiskers) and median lines are shown for $n = 4$ individuals per species and treatment.

The range of soil CH₄ fluxes under individuals of *Heisteria* (-190 - 539 µg m⁻² hr⁻¹) was greater than under individuals of *Simarouba* (-89.4 - 450 µg m⁻² hr⁻¹). Consequently, the mean soil CH₄ flux beneath *Heisteria* individuals was slightly more negative than that beneath *Simarouba* (-2.8 ±5.1 µg m⁻² hr⁻¹ and -2.3 ±5.5 µg m⁻² hr⁻¹ respectively).

Species and treatment effects on tree stem fluxes of CH₄

Surprisingly, tree stem CH₄ fluxes were mostly positive throughout the study period, even though the upland soils at the study site tended to act as a sink for methane during the dry season (Fig. 2). Accordingly, we found no relationship between soil and stem CH₄ fluxes. There was a significant species × treatment interaction on stem CH₄ fluxes, whereby *Heisteria* stems had higher CH₄ fluxes in L+ plots and lower stem fluxes in L- plots compared to *Simarouba* stems ($p < 0.001$, $\chi^2 = 24.5$; Fig. 4). Overall, the median CH₄ flux was very similar between the two species, with 72.6 µg m⁻² hr⁻¹ and 75.1 µg m⁻² hr⁻¹ for *Heisteria* and *Simarouba* respectively. Tree stem CH₄ fluxes in individuals of *Heisteria* were mostly positive, with a mean flux of 101 ±14.9 µg m⁻² hr⁻¹ over the dry-wet season transition, whereas the mean flux for *Simarouba* was lower at 87.7 ±18.5 µg m⁻² hr⁻¹.

Stem CH₄ fluxes in *Simarouba* displayed greater inter-week variability across all treatments (Fig. 4) and had a greater range throughout the study (-276 µg m⁻² hr⁻¹ to 678 µg m⁻² hr⁻¹) than those from *Heisteria* stems. CH₄ stem fluxes from *Heisteria* ranged from -156 to 598 µg m⁻² hr⁻¹ and remained relatively constant throughout the study in CT and L- plots but were much more variable in L+ plots (Fig. 4).




Figure 4. Methane (CH₄) fluxes from tree stems of *Heisteria concinna* (top panels) and *Simarouba amara* (bottom panels) in experimental litter removal (L-), control (CT) and litter addition (L+) treatments in a lowland tropical forest on upland soil in Panama, Central America; fluxes were measured weekly at 0.3-m height during the transition from the dry season (weeks 14-19) to the wet season (weeks 20-30); ranges (boxes and whiskers) and median lines are shown for $n = 4$ individuals per species and treatment.

269 Seasonal variation in N₂O fluxes

270 There was no clear seasonal pattern in soil or tree stem N₂O fluxes during the wet season
 271 (Fig. 5) and no effect of soil water content. However, we were unable to quantify N₂O fluxes
 272 during weeks 14-19, indicating very limited N₂O production during the dry season.




Figure 5 Nitrous oxide (N_2O) fluxes from tree stems at 0.3-m height (top panel) and soil (centre panel) in a lowland tropical forest on upland soil in Panama, Central America; means and standard errors are given for $n = 4$; measurements were made weekly during wet season (bottom panel) showing total rainfall (bars) and temperature (line).

275 There was a marginally significant additive effect of species and litter treatment for soil N₂O
 276 fluxes ($p = 0.092$, $\chi^2 = 6.45$), whereby soil N₂O fluxes measured beneath *Heisteria* individuals
 277 were greater in L+ plots compared to L- plots and tended to be higher than soil N₂O fluxes
 278 measured beneath *Simarouba* (means of $138 \pm 21.7 \mu\text{g m}^{-2} \text{h}^{-1}$ and $114 \pm 15.7 \mu\text{g m}^{-2} \text{h}^{-1}$,
 279 respectively; Fig. 6). Wet season soil N₂O fluxes beneath individuals of *Heisteria* ranged from
 280 -190 to $539 \mu\text{g m}^{-2} \text{h}^{-1}$ (median: $110 \mu\text{g m}^{-2} \text{h}^{-1}$) and N₂O fluxes under *Simarouba* trees ranged
 281 from -89.4 to $450 \mu\text{g m}^{-2} \text{h}^{-1}$ (median: $87.9 \mu\text{g m}^{-2} \text{h}^{-1}$).




Figure 6. Soil nitrous oxide (N₂O) fluxes under individuals of *Heisteria concinna* (top panels) and *Simarouba amara* (bottom panels) in experimental litter removal (L-), control (CT) and litter addition (L+) treatments in a lowland tropical forest on upland soil in Panama, Central America, measured weekly during the wet season; ranges (boxes and whiskers) and median lines are shown for $n = 4$ individuals per species and treatment.

282 *Species and treatment effects on tree stem fluxes of N₂O*

283 There was a significant additive effect of species and litter treatment on stem N₂O fluxes (χ^2
 284 = 9.66, $p = 0.022$), whereby fluxes from *Simarouba* stems were greater than those from
 285 *Heisteria* in L+ and CT plots, but not in L- plots (Fig. 7). Overall, median N₂O fluxes from
 286 *Heisteria* were much lower than those from *Simarouba* (101 $\mu\text{g m}^{-2} \text{h}^{-1}$ and 1001 $\mu\text{g m}^{-2} \text{h}^{-1}$,
 287 respectively) over the course of the study. N₂O fluxes from *Heisteria* stems were less variable
 288 (range: -2857 to 4270 $\mu\text{g m}^{-2} \text{h}^{-1}$) than *Simarouba* (range: -3770 to 8361 $\mu\text{g m}^{-2} \text{h}^{-1}$). Overall, a
 289 greater proportion of stem fluxes in *Simarouba* were positive and the mean stem flux from
 290 *Heisteria* was $80 \pm 234 \mu\text{g m}^{-2} \text{h}^{-1}$ compared to $1193 \pm 361 \mu\text{g m}^{-2} \text{h}^{-1}$ for *Simarouba* (Fig. 7).




Figure 7. Nitrous oxide (N₂O) fluxes from tree stems of *Heisteria concinna* (top panels) and *Simarouba amara* (bottom panels) in experimental litter removal (L-), control (CT) and litter addition (L+) treatments in a lowland tropical forest on upland soil in Panama, Central America; fluxes were measured weekly at 0.3-m height during the wet season (weeks 20-30); ranges (boxes and whiskers) and median lines are shown for $n = 4$ individuals per species and treatment.

Discussion

We demonstrate that upland soils in this seasonal tropical forest represented a sink for CH_4 during the dry season but became a source of CH_4 and N_2O during the transition to the wet season. Importantly, tree stems consistently emitted CH_4 during both the wet and dry seasons, which may offset the dry season soil sink. Tree stem fluxes of N_2O were only detectable in the wet season but were also mostly positive, indicating that tropical tree stems on upland soils could be a hitherto unaccounted for source of N_2O .

Seasonal patterns of soil CH_4 and N_2O fluxes

Soil trace GHG fluxes were strongly seasonal. The soil acted as a CH_4 sink during the dry season, when upper horizons of the soil profile have low soil water content and are well aerated and methanogenic archaea are probably dormant. With the onset of the wet season at the beginning of May (week 20; Fig. 2b), soil CH_4 fluxes shifted from strongly negative towards positive values, i.e. net CH_4 emissions. The linear relationship between soil CH_4 emissions and soil water content (Fig. 1) is typical of tropical and subtropical swamp forest and rainforest sites (Yu et al., 2008; Rowlings et al., 2012; Hall et al., 2013), and indicates that the higher soil water content during the wet season at our site created conditions that were more favourable for methanogenesis but less so for methanotrophy.

N_2O emissions usually increase with soil water content as the more productive anaerobic process of denitrification becomes dominant (Keller and Reiners, 1994) but we observed no relationship between soil water content and soil N_2O emissions in our study. However, we were unable to statistically assess seasonal patterns in N_2O fluxes because fluxes were not quantifiable for the majority of dry season measurements from the soil or tree stem bases. The lack of measurable N_2O production during the dry season at our study site is nonetheless

consistent with strong seasonality in N₂O fluxes, as the low variability in soil water content during the wet season explains why there was no relationship with N₂O. Alternatively, the timing and frequency of measurements may have been insufficient to detect relationships between soil water content and N₂O fluxes, as soil water saturation at our study site occurred as a result of heavy rainfall, rather than rising water table depth or flooding, and mesocosm studies have demonstrated that N₂O emissions were greatest 24 hours (Machacova et al. 2013) or ~45 hours (Liengaard et al. 2014) after rewetting but then declined rapidly.

We focussed primarily on soil water content as a control of trace GHG fluxes because temperature differences have a more profound effect on trace GHG fluxes when temperatures drop below 15°C (Castro *et al.*, 1995), and soil temperature was consistently >24°C in our study. Other factors such as diffusion rate (affected by e.g. soil density; Le Mer and Roger, 2001; Liu et al., 2007) and drought effects (Davidson et al., 2008; Itoh et al., 2010) become relatively more important at higher temperatures. It is also worth noting that the 2014 El Niño event resulted in a strong dry season in Panama, with lower rainfall than average, and seasonal effects in the present study may therefore have been exaggerated by the more severe dry season compared with other years.

Trace GHG fluxes from tree stems

Despite substantial changes in soil trace GHG fluxes between the dry and the wet season, there was no clear seasonal pattern in trace GHG fluxes from tree stems and therefore no support for our hypothesis of a relationship between CH₄ and N₂O fluxes from the soil and adjacent tree stems. Emissions of CH₄ and N₂O from tree stems in floodplain and wetland systems reflect the composition of soil trace GHG concentrations (Terazawa et al., 2007; Purvaja et al., 2004; Pangala et al., 2013) because the majority of CH₄ and N₂O emitted via the plant pathway originates from methanogenic consortia and denitrifying communities in the soil. However,

variation in tree water uptake and trace GHG production with soil depth could explain why we found no relationship between soil and tree stem CH₄ or N₂O fluxes. Previous work at the study site demonstrated that soil water content and N₂O concentrations increase with depth, whereas CH₄ concentrations were highest at c. 20-cm depth and then decreased with soil depth to c. 1.25-m (Koehler et al., 2012). Accordingly, we would expect higher CH₄ fluxes from stems when trees source water from shallower soil horizons, and higher N₂O fluxes when trees access water from deeper in the soil profile. Hence, the relationship between soil and tree stem GHG fluxes is likely to be influenced by the location of CH₄ or N₂O production in the soil profile, the rooting architecture of the tree species, and the preferential water uptake patterns of individual trees. Further, there is increasing evidence that CH₄ is also produced within tree stems (Wang et al., 2016) and methanogens can predominate in heartwood microbial communities (Yip et al., 2018), which could also contribute to CH₄ emissions from tree stems even when the soils were acting as a sink.

The majority of CH₄ and N₂O fluxes from tree stems were positive throughout the study (Figs. 4 & 7) and, unlike soil trace GHG fluxes, the rates did not change significantly between seasons. Consistent trace GHG emissions from tree stem bases has also been observed in temperate upland forests (Wang et al. 2016; Pitz & Megonigal, 2017; Warner et al., 2017) and suggests that GHGs may be produced within the tree stem (Covey et al. 2012; Wang et al. 2016), or that the transport of trace GHGs through tree stems bypassed the oxygenated surface horizons, where the majority of CH₄ oxidation (Teh et al., 2005; Wolf et al., 2012) and more complete denitrification (Koehler et al., 2012; Wieder et al., 2011) occurs. Hence, the production or transport of trace GHGs in tree stems could represent a large and currently unaccounted for source of N₂O and CH₄ emissions from tropical forests on upland soils. The median tree stem fluxes of CH₄ (~ 74 µg m⁻² h⁻¹) and N₂O (~99 µg m⁻² h⁻¹) we measured during the transition from the dry to the rainy season are comparable to fluxes measured in temperate

upland trees (Machacova et al., 2013; Wang et al., 2016; Wen et al., 2017; Pitz and Megonigal 2018) and our tree stem CH₄ fluxes also lie within the range of tropical peatland forests (17- 185 $\mu\text{g m}^{-2} \text{h}^{-1}$; Pangala et al., 2013).

We measured trace GHG fluxes from the stems of two common tree species with distinct life history strategies and wood densities because stem GHG fluxes are also likely to vary by species' physiological traits (Pangala et al., 2013). In our study, both CH₄ and N₂O fluxes were generally higher from stems of the fast-growing pioneer *Simarouba* (Fig. 4 & 7) except in the litter addition plots, where stem CH₄ fluxes were greater from the slow-growing shade-tolerant tree *Heisteria*. Higher stem emissions from the canopy species *Simarouba* could possibly be explained by greater rates of evapotranspiration and lower wood density compared to the subcanopy species *Heisteria*. Mesocosm studies of two temperate tree species common to forests on upland soils found that fast-growing species created channels of greater gas diffusivity because they had higher fine root density and greater maximum root depth (Fender et al., 2013). Further, higher rates of net primary productivity in canopy species may result in greater root exudation, which could stimulate CH₄ and N₂O production (Butterbach-Bahl et al., 2002). The potential influence of different tree species traits on stem trace GHG fluxes makes it challenging to assess ecosystem-level fluxes in highly diverse tropical forests, especially as CH₄ and N₂O concentrations within soil pore gas and water are also influenced tree diversity (Machacova et al., 2013; Warner et al., 2017). However, we demonstrate that tree species with distinct life history strategies and ecological niches emit CH₄ and N₂O, from their stems, which suggests that trace GHG fluxes from tree stems could be widespread in tropical forests on upland soils.

Interestingly, tree stems also acted as a sink for trace GHG gases in this study (Figs. 4 and 7). It is conceivable that changes in CH₄ and N₂O concentrations in soil water at different depths could generate a diffusion gradient from the atmosphere into tree stems, thus resulting in tree

stem uptake of trace GHGs, possibly as a result of active consumption by epiphytic and endophytic methanotrophs. This intriguing possibility merits further attention.

Links between trace GHG fluxes, seasonality, and decomposition processes

The decomposition of plant litter plays a role in methanogenesis and nitrous oxide emissions because it releases the labile carbon that supplies electron donors for methanogens (Megonigal and Guenther 2008) and the nitrate for denitrification (Teh et al., 2008). In upland tropical forest soils, the combined spatial variability in soil water content, decomposition processes, and electron donors may represent a greater control on trace GHG uptake or emission than any single factor. At our study site, a large proportion of the annual litterfall occurs during the dry season (Sayer & Tanner 2010), and the rapid increase in soil fluxes of CH₄ and N₂O at the start of the wet season are likely to result from a combination of increased water-filled pore space and the decomposition of the thick litter that accumulates during the dry season (Wieder and Wright, 1995). Enhanced decomposition immediately after wetting (Vasconcelos *et al.*, 2007) could lead to spikes in N₂O fluxes that coincide with rainfall (e.g. weeks 24 and 28 in Fig. 5). Correspondingly, the low rates of decomposition could also partly explain why we were unable to detect N₂O fluxes in the majority of dry season measurements.

Given the influence of decomposition processes on trace GHG production, we hypothesised that both CH₄ and N₂O emissions would be higher in litter addition plots as a result of greater availability in substrate for soil microorganisms. Although there were no consistent effects of the litter treatments on trace GHG emissions, higher N₂O emissions in litter addition plots towards the end of the study period (weeks 24-30; Fig. 6 & 7) and the additive or interactive effect of litter addition and species on tree stem N₂O and CH₄ fluxes (Fig. 4) suggest that there is a connection between trace GHG fluxes and litter quantity. However, increased litter inputs are only likely to result in increased N₂O emissions where there are sufficiently large

communities of denitrifying bacteria that can respond quickly to wetting events, creating hotspots and 'hot moments' (McClain et al. 2003). The high inter-week variation in tree stem N_2O fluxes in the present study (Fig. 7) could therefore also be explained by the timing and presence of such hotspots within the rooting zone.

Overall, the discrepancies between soil and stem trace GHG fluxes, and the apparent stronger effect of litter manipulation on fluxes of CH_4 and N_2O from *Heisteria* tree stems compared to *Simarouba* stems (Fig. 4 and 7) may be attributed to differences in species' rooting depths and the gradients of trace GHG production with soil depth. The possible role of rooting depth as a control of trace GHG fluxes from tree stems on upland soils is another avenue of investigation that requires further attention.

3.5 Conclusions

Our results suggest that tropical tree stems on upland soils may represent a hitherto unaccounted for conduit of trace GHG emissions, which likely originate from deeper soil horizons. Together, litter addition and tree species identity influenced stem CH_4 and N_2O fluxes, which suggests that tree uptake and emissions of trace GHGs is influenced both by decomposition processes and species traits. Given that tropical forests on upland soils cover a larger area than tropical wetland forests, the mechanisms underlying the fluxes of CH_4 and N_2O from tree stems, and their contribution to global trace GHG budgets, merits further examination.

Acknowledgements

We thank M. Peacock and P. Rafferty for assistance with laboratory analyses, L. Bréchet, F. Valdez and B Castro for help with fieldwork, and E.V.J. Tanner for discussions about the litter manipulation experiment; BW was supported by a studentship from the Natural Environment Research Council, UK; VG acknowledges support from the AXA Research Fund and NERC

grant NE/J010928/1; EJS was supported by funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013), ERC Grant Agreement No. 307888. The authors have no conflict of interest to declare.

References

- Alm, J., Shurpali, N.J., Tuittila, E.S., Laurila, T., Maljanen, M., Saarnio, S. & Minkkinen, K. (2007). Methods for Determining Emission Factors for the Use of Peat and Peatlands - Flux Measurements and Modelling. *Boreal Environment Research*, 12:85-100.
- Baird, A.J., Stamp, I., Heppell, C.M. & Green, S.M. (2010). CH₄ flux from peatlands: A new measurement method. *Ecohydrology*, 3, 360-367.
- Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed Effects Models Using lme4. *Journal of Statistical Software*, 67, 1-48.
- Bouwman, A. V., Van der Hoek, K. W., & Olivier, J. G. J. (1995). Uncertainties in the global source distribution of nitrous oxide. *Journal of Geophysical Research: Atmospheres*, 100, 2785-2800.
- Butterbach-Bahl, K. & Papen, H. (2002). Four years continuous record of CH₄-exchange between the atmosphere and untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany. *Plant and Soil*, 240, 77-90.
- Carmichael, M.J., Bernhardt, E.S., Bräuer, S.L. & Smith, W.K. (2014). The role of vegetation in methane flux to the atmosphere: Should vegetation be included as a distinct category in the global methane budget? *Biogeochemistry*, 119, 1-24.
- Castro, M.S., Steudler, P.A., Melillo, J.M., Aber, J.D. & Bowden, R.D. (1995). Factors controlling atmospheric methane consumption by temperate forest soils. *Global Biogeochemical Cycles*, 9, 1-10.
- Cavelier, J. (1992). Fine-root biomass and soil properties in a semideciduous and a lower

- montane rain forest in Panama. *Plant and Soil*, 142, 187-201.
- Condit, R., Windsor, D.M. & Hubbell, S.P. (2013). NPP Tropical Forest: Barro Colorado, Panama, 1969-2000, R1. Data set. Available at: <http://daac.ornl.gov>.
- Cooper, M.D., Evans, C.D., Zielinski, P., Levy, P.E., Gray, A., Peacock, M., Norris, D., Fenner, N. & Freeman, C. (2014). Infilled ditches are hotspots of landscape methane flux following peatland re-wetting. *Ecosystems*, 17, 1227-1241.
- Covey, K.R., Wood, S.A., Warren, R.J., Lee, X. and Bradford, M.A. (2012). Elevated methane concentrations in trees of an upland forest. *Geophysical Research Letters*, 39, 15.
- Davidson, E.A., Nepstad, D.C., Ishida, F.Y. & Brando, P.M (2008). Effects of an experimental drought and recovery on soil emissions of carbon dioxide, methane, nitrous oxide, and nitric oxide in a moist tropical forest. *Global Change Biology*, 14, 2582-2590.
- Fender, A.C., Gansert, D., Jungkunst, H.F., Fiedler, S., Beyer, F., Schützenmeister, K., Thiele, B., Valtanen, K., Polle, A. & Leuschner, C. (2013). Root-induced tree species effects on the source/sink strength for greenhouse gases (CH₄, N₂O and CO₂) of a temperate deciduous forest soil. *Soil Biology and Biochemistry*, 57, 587-597.
- Gauci, V., Gowing, D.J., Hornibrook, E.R., Davis, J.M. & Dise, N.B. (2010). Woody stem methane emission in mature wetland alder trees. *Atmospheric Environment*, 44, 2157-2160.
- Graffmann, K., Grosse, W., Junk, W.J. & Parolin, P. (2008). Pressurized gas transport in Amazonian floodplain trees. *Environmental and Experimental Botany*, 62, 371–375
- Hall, S.J., McDowell, W.H. & Silver, W.L. (2013). When wet gets wetter: Decoupling of moisture, redox biogeochemistry, and greenhouse gas fluxes in a humid tropical forest soil. *Ecosystems*, 16, 576-589.
- Houghton, J.T., Ding, Y., Griggs, D.J., Noguer, M., Van Der Linden, P.J., Dai, X., Maskell,

- K. & Johnson, C.A. (2001). *Climate change 2001: the scientific basis. Contribution of working group I to the third assessment report of the International Panel on Climate Change*. Cambridge University Press, UK, 36, p.44.
- IPCC (2007). *Climate Change 2007 - The Physical Science Basis: Working Group I Contribution to the Fourth Assessment Report of the IPCC*, Cambridge University Press, Cambridge, UK and New York, USA.
- IPCC (2013). *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, USA.
- Itoh, M., Kosugi, Y., Takanashi, S., Hayashi, Y., Kanemitsu, S., Osaka, K., Tani, M. & Nik, A.R. (2010). Temporal and spatial variations of soil carbon dioxide, methane, and nitrous oxide fluxes in a Southeast Asian tropical rainforest. *Biogeosciences Discussions*, 7, 6847-6887.
- Keller, M. & Reiners, W.A. (1994). Soil-atmosphere exchange of nitrous oxide, nitric oxide, and methane under secondary succession of pasture to forest in the Atlantic lowlands of Costa Rica. *Global Biogeochemical Cycles*, 8, 399-404.
- Koehler, B., Corre, M.D., Steger, K., Well, R., Zehe, E., Sueta, J.P. & Veldkamp, E. (2012). An in-depth look into a tropical lowland forest soil: Nitrogen-addition effects on the contents of N₂O, CO₂ and CH₄ and N₂O isotopic signatures down to 2-m depth. *Biogeochemistry*, 111, 695-713.
- Kreuzwieser, J., Buchholz, J. & Rennenberg, H. (2003). Emission of methane and nitrous oxide by Australian mangrove ecosystems. *Plant Biology*, 5, 423-431.
- Krithika, K., Purvaja, R. & Ramesh, R. (2008). Fluxes of methane and nitrous oxide from an Indian mangrove. *Current Science*, 94, 218-224.
- Le Mer, J. & Roger, P. (2001). Production, oxidation, emission and consumption of methane

- by soils: a review. *European Journal of Soil Biology*, 37, 25-50.
- Leigh, E.G. (1999). *Tropical Forest Ecology : A View from Barro Colorado Island*. Oxford University Press, Oxford, UK.
- Li, C.S., Aber, J., Stange, F., Butterbach-Bahl, K. & Papen, H. (2000), A process-oriented model of N₂O and NO emissions from forest soils: 1. Model development, *Journal of Geophysical Research*, 105, 4369-4384.
- Liengaard, L., Figueiredo, V., Markfoged, R., Revsbech, N.P., Nielsen, L.P., Prast, A.E. & Kühl, M. (2014). Hot moments of N₂O transformation and emission in tropical soils from the Pantanal and the Amazon (Brazil). *Soil Biology and Biochemistry*, 75, 26-36.
- Liu, C., Holst, J., Brüggemann, N., Butterbach-Bahl, K., Yao, Z., Yue, J., Han, S., Han, X., Krümmelbein, J., Horn, R. & Zheng, X. (2007). Winter-grazing reduces methane uptake by soils of a typical semi-arid steppe in Inner Mongolia, China. *Atmospheric Environment*, 41, 5948-5958.
- Machacova, K., Papen, H., Kreuzwieser, J. & Rennenberg, H. (2013). Inundation strongly stimulates nitrous oxide emissions from stems of the upland tree *Fagus sylvatica* and the riparian tree *Alnus glutinosa*. *Plant and Soil*, 364, 287-301.
- Machacova, K., Bäck, J., Vanhatalo, A., Halmeenmäki, E., Kolari, P., Mammarella, I., Pumpanen, J., Acosta, M., Urban, O. & Pihlatie, M. (2016). *Pinus sylvestris* as a missing source of nitrous oxide and methane in boreal forest. *Scientific Reports*, 6, 23410.
- McClain, M.E., Boyer, E.W., Dent, C.L., Gergel, S.E., Grimm, N.B., Groffman, P.M., Hart, S.C., Harvey, J.W., Johnston, C.A., Mayorga, E. & McDowell, W.H. (2003). Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic ecosystems. *Ecosystems*, 6, 301-312.
- Megonigal, J.P. & Guenther, A.B. (2008). Methane emissions from upland forest soils and

540 vegetation. *Tree Physiology*, 28, 491-498.

541 Pan, Y., Birdsey, R.A., Phillips, O.L. & Jackson, R.B. (2013). The structure, distribution, and
542 biomass of the world's forests. *Annual Review of Ecology, Evolution, and Systematics*, 44,
543 593-622

544 Pangala, S.R., Moore, S., Hornibrook, E.R. & Gauci, V. (2013). Trees are major conduits for
545 methane egress from tropical forested wetlands. *New Phytologist*, 197, 524-531.

546 Pangala, S.R., Gowing, D.J., Hornibrook, E.R. & Gauci, V. (2014). Controls on methane
547 emissions from *Alnus glutinosa* saplings. *New Phytologist*, 201: 887-896.

548 Pangala, S.R., Enrich-Prast, A., Basso, L.S., Peixoto, R.B., Bastviken, D., Hornibrook, E.R.,
549 Gatti, L.V., Marotta, H., Calazans, L.S.B., Sakuragui, C.M., Bastos, W.R., Malm, O.,
550 Gloor, E., Bharat Miller, J. & Gauci, V. (2017). Large emissions from floodplain trees
551 close the Amazon methane budget. *Nature*, 552, 230.

552 Pinheiro, J.C. & Bates, D.M., 2000. (2000) Linear Mixed-Effects Models: Basic Concepts
553 and Examples. In: *Mixed-Effects Models in S and S-PLUS*. Statistics and Computing.
554 Springer, New York, NY, pp.3-56.

555 Pitz, S. and Megonigal, J.P. (2017). Temperate forest methane sink diminished by tree
556 emissions. *New Phytologist*, 214, 1432-1439.

557 Pitz, S. L., Megonigal, J. P., Chang, C. H., & Szlavecz, K. (2018). Methane fluxes from tree
558 stems and soils along a habitat gradient. *Biogeochemistry*, 137, 307-320.

559 Purvaja, R., Ramesh, R. & Frenzel, P. (2004). Plant-mediated methane emission from an
560 Indian mangrove. *Global Change Biology*, 10, 1825-1834.

561 R Core Team (2016). *R: A Language and Environment for Statistical Computing*. R
562 Foundation for Statistical Computing, Vienna Austria.

563 Rice, A.L., Butenhoff, C.L., Shearer, M.J., Teama, D., Rosenstiel, T.N. & Khalil, M.A.K.
564 (2010). Emissions of anaerobically produced methane by trees. *Geophysical Research*

Letters, 37, 1-6.

Rowlings, D.W., Grace, P.R., Kiese, R. & Weier, K.L. (2012). Environmental factors controlling temporal and spatial variability in the soil-atmosphere exchange of CO₂, CH₄ and N₂O from an Australian subtropical rainforest. *Global Change Biology*, 18, 726-738.

Rusch, H. & Rennenberg, H. (1998). Black alder (*Alnus glutinosa* (L.) Gaertn.) trees mediate methane and nitrous oxide emission from the soil to the atmosphere. *Plant and Soil*, 201, 1-7.

Sayer, E.J., Tanner, E.V.J. & Cheesman, A.W. (2006). Increased litterfall changes fine root distribution in a moist tropical forest. *Plant and Soil*, 281, 5-13.

Sayer, E.J. & Tanner, E.V.J. (2010). Experimental investigation of the importance of litterfall in lowland semi-evergreen tropical forest nutrient cycling. *Journal of Ecology*, 98, 1052-1062.

Siegenthaler, A., Welch, B., Pangala, S. R., Peacock, M., & Gauci, V. (2016) Technical Note: Semi-rigid chambers for methane gas flux measurements on tree stems, *Biogeosciences*, 13, 1197-1207.

Tang, X., Liu, S., Zhou, G., Zhang, D. & Zhou, C. (2006). Soil-atmospheric exchange of CO₂, CH₄, and N₂O in three subtropical forest ecosystems in southern China. *Global Change Biology*, 12, 546-560.

Teh, Y.A., Dubinsky, E.A., Silver, W.L. & Carlson, C.M. (2008). Suppression of methanogenesis by dissimilatory Fe(III)- reducing bacteria in tropical rain forest soils: Implications for ecosystem methane flux. *Global Change Biology*, 14, 413-422.

Teh, Y.A., Silver, W.L. & Conrad, M.E. (2005). Oxygen effects on methane production and oxidation in humid tropical forest soils. *Global Change Biology*, 11, 1283-1297.

Terazawa, K., Ishizuka, S., Sakata, T., Yamada, K. & Takahashi, M. (2007). Methane emissions from stems of *Fraxinus mandshurica* var. *japonica* trees in a floodplain forest.

Soil Biology and Biochemistry, 39, 2689-2692.

Vasconcelos, S.S., Zarin, D.J., Da Rosa, M.B.S., de Assis Oliveira, F. & De Carvalho, C.J.R.

(2007). Leaf decomposition in a dry season irrigation experiment in eastern Amazonian

forest regrowth. *Biotropica*, 39, 593-600.

Wang, Z.P., Gu, Q., Deng, F.D., Huang, J.H., Megonigal, J.P., Yu, Q., Lü, X.T., Li, L.H.,

Chang, S., Zhang, Y.H. & Feng, J.C. (2016). Methane emissions from the trunks of living

trees on upland soils. *New Phytologist*, 211, 429-439.

Warner, D.L., Villarreal, S., McWilliams, K., Inamdar, S. & Vargas, R. (2017). Carbon

dioxide and methane fluxes from tree stems, coarse woody debris, and soils in an upland

temperate forest. *Ecosystems*, 1-12.

Wen, Y., Corre, M.D., Rachow, C., Chen, L. and Veldkamp, E. (2017). Nitrous oxide

emissions from stems of alder, beech and spruce in a temperate forest. *Plant and Soil*, 420,

423-434.

Wieder, R.K. & Wright, S.J. (1995). Tropical forest litter dynamics and dry season irrigation

on Barro Colorado Island, Panama. *Ecology*, 76, 1971-1979.

Wieder, W.R., Cleveland, C.C. & Townsend, A.R. (2011). Throughfall exclusion and leaf

litter addition drive higher rates of soil nitrous oxide emissions from a lowland wet tropical

forest. *Global Change Biology*, 17, 3195-3207.

Wolf, K., Flessa, H. & Veldkamp, E. (2012). Atmospheric methane uptake by tropical

montane forest soils and the contribution of organic layers. *Biogeochemistry*, 111, 469-483

Yip, D. Z., Veach, A. M., Yang, Z. K., Cregger, M. A., & Schadt, C. W. (2018). Methanogenic

Archaea dominate mature heartwood habitats of Eastern Cottonwood (*Populus*

deltoides). *New Phytologist*, doi: 10.1111/nph.15346.

Yu, K., Faulkner, S.P. & Baldwin, M.J. (2008). Effect of hydrological conditions on nitrous

oxide, methane, and carbon dioxide dynamics in a bottomland hardwood forest and its

615 implication for soil carbon sequestration. *Global Change Biology*, 14, 798-812.
616 Zhuang, Q., Lu, Y. & Chen, M. (2012). An inventory of global N₂O emissions from the soils
617 of natural terrestrial ecosystems. *Atmospheric Environment*, 47, 66-75.
618

ACCEPTED VERSION